A Long Quest Towards Improved GHG Inventories

Guido Franco
Public Interest Energy Research (PIER) Program
California Energy Commission

Jorn Herner
Research Division
California Air Resources Board

Sixth California Annual Climate Change Research Symposium Sacramento, California September 8 -10, 2009



Outline

- Part I [Guido Franco PIER/CEC]
 - How accurate are the existing GHG inventories?
 Some observations
 - PIER supported research projects on GHG inventory methods in California
- Part II [Jorn Herner ARB]
 - Improving Greenhouse Gas Inventories and Validation Field Studies

Disclaimer

The views and statements including in this presentation do not necessarily represent the views and opinions of the Energy Commission, Air Resources Board, or the State of California




How accurate are the existing GHG inventories? Some observations



CO₂ from the Combustion of Fossil Fuels in the US

- Baseline (1990) US
 emission estimates
 reported by the US EPA
 have gone down since their
 first inventory released in
 1998
- Changes are relatively minor (~ 3%)
- Reported uncertainty -2% to 6%*
- What is the situation with State level data?

Normalized **1990** CO₂ Emissions from Fossil Fuel Combustion – Different Annual US EPA Reports




Source: Franco and Oliver, 2009. Data from different US EPA annual inventory reports




Comparing EIA and ARB Estimates for CA

- In general the CO₂ emission estimates from EIA and ARB are relatively close.
- ARB used the CALEB data set prepared for PIER by LBNL but ARB enhanced CALEB for the inventory.
- Comparing CALEB with secondary sources suggests that the largest sources of uncertainty* are:
 - differences between supply and consumption data (-2% to +4%)
 - carbon emission factor uncertainties (-1% to +5%)

Relative Differences of CO₂ Emissions from the Combustion of Fossil Fuels in California: EIA vs ARB






* Source: LBNL 2008



Non-CO₂ gases


- IPCC/US EPA methods are highly uncertain
- Estimated US 1990
 baseline year N₂O
 emissions from the
 application of fertilizer
 have changed
 substantially
- Estimate of CH₄
 emissions from landfills
 are stabilizing but......


Normalized **1990** CH₄ from Landfills and N₂O Emissions From Fertilizer Applications



Modeled Methane Landfill Generation* is not a Very Good Predictor of Emissions

- Current Inventory Method
 emissions = (generation recovery) x (1 OF)
- Assumed cover soil oxidation fraction (OF) = 10%
- Estimated error is about -37% to 18% **
- However, the error can be much larger if there are fundamental structural problems with the method used to estimate emissions





^{*}using a multi-component first order kinetic model and detailed waste input data
** 2009 Draft U.S. Greenhouse Gas Inventory Report. US EPA. February 2009


Source: Jean Bogner. TAC meeting March 11, 2009 Spokas et al. 2006




Methane Oxidation Rates as a Function of Moisture and Temperature

(Marina Landfill, CA)

- Oxidation is a function of multiple factors including moisture, temperature, and type of cover
- A new model (IPPC Tier 4) is being developed for CA





Soil incubation experiments

Source: Jean Bogner. TAC Meeting. March 11, 2009




PIER Supported Research Projects in California



GHG Inventory Methods: PIER Projects

- California Energy Balances:
 Phase II
- Non-CO₂ gases:
 - Methane emissions from landfills (IWMB, ARB)
 - GHG emissions from dairy farms (CDFA, ARB)
 - N₂O emissions from fertilizers (ARB, CDFA)
 - Net GHG emissions from C sequestration in terrestrial ecosystems (DWR, USGS, ARB)
- Measurement of GHG in ambient air (ARB, NOAA)
- Fugitive methane emissions from the natural gas system (ARB)







Relevant PIER Reports Research Powers the Future

- Observation of CH₄ and other Non-CO₂ Green House Gas Emissions from California (Forthcoming)
- Discussion paper on GHG inventory methods: research needs. September 2008.
- Developing and Applying Process-Based Model for Estimating Greenhouse Gas and Air Emissions From California Dairies. Feb 2009
- Assessment of Greenhouse Gas Mitigation in California Agricultural Soils. Jan 2009.
- Summary of Rangelands A Suitable for Terrestrial Carbon Sequestration In Shasta County. Feb 2008.
- Carbon Supply from Changes in Management of Forest and Rangelands in Shasta County, California, June 2007.
- Assessing Impacts of Rangeland Management and Reforestation of Rangelands on Greenhouse Gas Emissions: A Pilot Study for Shasta County. Feb 2007.
- Carbon Supply from Changes in Management of Forest, Range, and Agricultural Lands in California: Forest Fuel Reduction. October 2006.
- Development of an Implementation Plan for Atmospheric Carbon Monitoring in California. August 2005.
- Research Roadmap for Greenhouse Gas Inventory Methods. July 2005
- Development of Energy Balances for the State of California. June 2005.
- Baseline Greenhouse Gas Emissions for Forest, Range, and Agricultural Lands In California.
 December 2004.
- Quantifying Carbon Dynamics and Greenhouse Gas Emissions in Agricultural Soils of California:
 A Scoping Study. October 2004

http://www.energy.ca.gov/publications/searchReports.php?pier1=Climate%20Change



- Improving GHG Inventory methods will require well funded and well coordinated research programs at the local, state, and federal levels. What ARB and CEC are doing is a good starting point.
- Every state agency with expertise/responsibility for different sources must be involved (e.g., CalFire and carbon stocks in forest)
- Long-term ambient monitoring must be continued and enhanced.
 Inventories with adequate temporal and spatial resolution are also needed.
- Maybe it is time to harmonize criteria and GHG inventories. There is a danger in trying to use existing models which may rely on indirect activity data (e.g., miles travelled) instead of more easily observable data (e.g., fuel consumption)



Improving Greenhouse Gas Inventories and Validation Field Studies

Jorn Dinh Herner
California Air Resources Board



Legislative Drivers

Assembly Bill 1493

Reduce GHG emissions from motor vehicles

Assembly Bill 32

 The State Board shall determine what the statewide greenhouse gas level was in 1990, and approve a statewide greenhouse gas emission limit that is equivalent to that level, to be achieved by 2020



California Governor Arnold Schwarzenegger signs AB32


Assembly Bill 1803

 Directs ARB to prepare, adopt, and update the California GHG Inventory



Purpose / Goal


- Inform the process to reach the AB32 Target
 - Support regulatory efforts
 - Assess the effectiveness of the mitigation program
 - Validate the GHG Inventory
 - Identify new sources / Improve emission factors
- Use a combination of in-house, collaborative, and extramural research






Strategy

- Apply same principles that has guided the last 20 years of research and improvements in vehicular emissions inventories to GHG inventory.
- Coordinated extramural, interagency, and in-house efforts
- Efforts on CO_2 as well as other GHGs, such as N_2O , CH_4 , HFCs, ODSs, and Black Carbon Nitrous Hydrofluorocarbons
 - Regional validation studies
 - Measure individual sources





Improving N₂O Emissions Estimates




- 1. N₂O Emissions from the Application of Fertilizers in Agricultural Soils (CEC)
- 2. Measuring and Modeling N₂O in CA Cotton, Corn and Vegetable Systems (CDFA)
- 3. How to Reduce N₂O Emissions from Nitrogen Land Application (ARB)



- The daily sea breeze pushes urban air against the base of the San Gabriel Mts. and solar heating provides vertical motion.
- The very steep, south-facing slope provides strong, rapid upslope flow of well-mixed air from the urban lowland mid-morning to early evening almost every day.
- Subsidence flow replaces the urban air with uncontaminated air from the lower free troposphere (LFT) at night, generally from the eastern North Pacific Ocean.

Mt Wilson - Los Angeles County Methane Emissions



 $\Delta CH_4 = 0.50 \text{ x } \Delta CO \text{ (based on correlation slope)}$

2007 LA County CO inventory 1,800 metric tons/day

 CH_4 emissions = (CO metric tons) x 0.50 x (16/28) = 4.0 MM tons CO_2 eq./year

Methods	LA County CH ₄ [MM Tons CO ₂ eq./year]	Notes
CH ₄ to CO Scaling Measured at Mt. Wilson	4.0 ± 0.35	Estimate
CARB CH ₄ inventory	3.0	IPCC methods

Establish a California Pilot Methane Network

- Provide greater spatial and temporal detail
- •Establish additional stations in California measuring CH₄, CO₂ and CO to get coverage of the Central Valley (ARB)
- Build on modeling work already funded by CEC
- Develop inverse models to validate methane inventory and find over or underestimated sources (extramural)





Measure Individual Sources

- •Develop the ability to measure GHG from individual sources:
 - •Two electric vehicles capable of measuring CO₂, CO, CH₄, N₂O, PM_{2.5}, BC, Temp, RH, Wind speed and direction, position (GPS).
 - •Use dispersion models, tracer gas release and flux chambers to make measurements





Measure Individual Sources



CalNex 2010

Research at the Nexus of Air Quality and Climate"

Combine the resources of NOAA and ARB in a major field study to achieve a step change in our understanding.

- •Some species affect both air quality and climate
- •Measurements not previously possible
- •Address the entire State and over the Pacific
- Answer foundational science issues





Thanks!

Guido Franco

Technical Lead for Climate Change Research
Public Interest Energy Research (PIER) Program
California Energy Commission
gfranco@energy.state.ca.us
(916)654-3940

Jorn Herner, Ph.D.

Manager, Greenhouse Gas Technology and Field Testing Section
Research Division
California Air Resources Board
jherner@arb.ca.gov
(916)324-9299



California Sixth Annual Climate Change Symposium

September 8 -10, 2009

Sacramento

Sacramento

Sacramento